

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 :
C12N 15/55, 9/20, C12Q 1/68, C12N
15/63, A61K 38/46, C07K 16/40

A2

(11) International Publication Number: WO 00/24911

(43) International Publication Date: 4 May 2000 (04.05.00)

(21) International Application Number: PCT/US99/25021

(22) International Filing Date: 27 October 1999 (27.10.99)

(30) Priority Data:

09/181,317	27 October 1998 (27.10.98)	US
Not furnished	27 October 1998 (27.10.98)	US
09/234,726	21 January 1999 (21.01.99)	US
Not furnished	21 January 1999 (21.01.99)	US

(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications

US	09/181,317 (CIP)
Filed on	27 October 1998 (27.10.98)
US	Not furnished (CIP)
Filed on	27 October 1998 (27.10.98)
US	09/234,726 (CIP)
Filed on	21 January 1999 (21.01.99)
US	Not furnished (CIP)
Filed on	21 January 1999 (21.01.99)

(71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive, #12, Mountain View, CA 94040 (US). BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). GUEGLER, Karl, J. [CH/US]; 1048 Oakland Avenue, Menlo Park, CA 94025 (US). CORLEY, Neil, C. [US/US]; 1240 Dale Avenue, #30, Mountain View, CA 94040 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont Park, San Jose, CA 95136 (US).

(74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

Without international search report and to be republished upon receipt of that report.

(54) Title: HUMAN PHOSPHOLIPASES

1	M E L A L L C G L V V M A	-	G V I P I Q G G I L N L N K M V	2641779
1	M K V L L L A V V I M A	F	G S I Q V Q G S L L E F G Q M I	GI 204319
30	K Q V T G K M P I L S Y W P Y G C H C G L G G R G Q P K D A			2641779
31	L F K T G K R A D V S Y G F Y G C H C G V G G R G S P K D A			GI 204319
60	T D W C C Q T H D C C Y D H L K T Q G C G I Y K D Y Y R Y N			2641779
61	T D W C C V T H D C C Y N R L E K R G C G T K F L T Y K F S			GI 204319
90	F S Q G N I H C S D K G S W C E Q Q L C A C D K E V A F C L			2641779
91	Y R G G Q I S C S T N Q D S C R K Q L C Q C D K A A A E C F			GI 204319
120	K R N L D T Y Q K R L R F Y W R P H C R G Q T P G C			2641779
121	A R N K K S Y S L K Y Q F Y L N K F C K G K T P S C			GI 204319

(57) Abstract

The invention provides human phospholipases (HPPL) and polynucleotides which identify and encode HPPL. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of HPPL.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
ER	Estonia						

HUMAN PHOSPHOLIPASES

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of human phospholipases and to the use of these sequences in the diagnosis, treatment, and prevention of cancer, autoimmune/inflammatory disorders, and reproductive disorders.

5

BACKGROUND OF THE INVENTION

Triglyceride lipases are enzymes that hydrolyze the ester bond of triglycerides. Lipases are widely distributed in animals, plants, and prokaryotes. In higher vertebrates there are at least three tissue-specific isozymes including pancreatic, hepatic, and gastric lipases. These three types of lipases are structurally closely related to each other as well as to lipoprotein lipase. Gastric lipases in the intestine aid in the digestion and absorption of dietary fats. Adipocytes contain lipases that break down stored triacylglycerols, releasing fatty acids for export to other tissues where they are required as fuel. Gastric, hepatic, and pancreatic lipases hydrolyze lipoprotein triglycerides and phospholipids. The most conserved region in pancreatic, hepatic, and gastric lipases is centered around a serine residue which is also present in lipases of prokaryotic origin.

Phospholipases, a group of enzymes that catalyze the hydrolysis of membrane phospholipids, are classified according to the bond cleaved in a phospholipid. They are classified into PLA1, PLA2, PLB, PLC, and PLD. Phospholipases are involved in many inflammatory reactions by making arachidonate available for eicosanoid biosynthesis. More specifically, arachidonic acid is processed into bioactive lipid mediators of inflammation such as lyso-platelet-activating factor and eicosanoids. The synthesis of arachidonic acid from membrane phospholipids is the rate-limiting step in the biosynthesis of the four major classes of eicosanoids which are prostaglandins, prostacyclins, thromboxanes and leukotrienes, and are involved in pain, fever, and inflammation. (Kaiser, E. et al. (1990) Clin. Biochem. 23:349-370.) Furthermore, leukotriene-B4 is known to function in a feedback loop which further increases PLA2 activity. (Wijkander, J. et al. (1995) J. Biol. Chem. 270:26543-26549.)

The secretory phospholipase A₂ (PLA2) superfamily comprises a number of heterogeneous enzymes whose common feature is to hydrolyze the *sn*-2 fatty acid acyl ester bond of phosphoglycerides. Hydrolysis of the glycerophospholipids releases free fatty acids and lysophospholipids. PLA2 activity generates precursors for the biosynthesis of biologically active lipids, hydroxy fatty acids, and platelet-activating factor.

PLA2s were first described as components of snake venoms, and were later characterized in numerous species. PLA2s have traditionally been classified into several major groups and subgroups

based on their amino acid sequences, divalent cation requirements, and location of disulfide bonds. The PLA2s of Groups I, II, and III consist of low molecular weight, secreted, Ca^{2+} -dependent proteins. Group IV PLA2s are primarily 85-kDa, Ca^{2+} -dependent cytosolic phospholipases. Finally, a number of Ca^{2+} -independent PLA2s have been described, which comprise Group V. (Davidson, F. F.

5 and Dennis, E. A., (1990) *J. Mol. Evol.* 31: 228-238; and Dennis, E. F. (1994) *J. Biol. Chem.* 269:13057-13060.)

The first PLA2s to be extensively characterized were the Group I, II, and III PLA2s found in snake and bee venoms. These venom PLA2s share many features with mammalian PLA2s including a common catalytic mechanism, the same Ca^{2+} requirement, and conserved primary and tertiary
10 structures. In addition to their role in the digestion of prey, the venom PLA2s display neurotoxic, myotoxic, anticoagulant, and proinflammatory effects in mammalian tissues. This diversity of pathophysiological effects is due to the presence of specific, high affinity receptors for these enzymes on various cells and tissues. (Lambeau, G. et al. (1995) *J. Biol. Chem.* 270:5534-5540.)

PLA2s from Groups I, IIA, IIC, and V have been described in mammalian and avian cells,
15 and were originally characterized by tissue distribution, although the distinction is no longer absolute. Thus, Group I PLA2s were found in the pancreas, Group IIA and IIC derived from inflammation-associated tissues (e.g., the synovium), and Group V were from cardiac tissue. The pancreatic PLA2s function in the digestion of dietary lipids and have been proposed to play a role in cell proliferation, smooth muscle contraction, and acute lung injury. The inflammatory PLA2s are potent mediators of
20 inflammatory processes and are highly expressed in serum and synovial fluids of patients with inflammatory disorders. These Group II PLA2s are found in most human cell types assayed and are expressed in diverse pathological processes such as septic shock, intestinal cancers, rheumatoid arthritis, and epidermal hyperplasia. A Group V PLA2 has been cloned from brain tissue and is strongly expressed in heart tissue. A human PLA2 was recently cloned from fetal lung, and based on
25 its structural properties, appears to be the first member of a new group of mammalian PLAs, referred to as Group X. Other PLA2s have been cloned from various human tissues and cell lines, suggesting a large diversity of PLA2s. (Chen, J. et al. (1994) *J. Biol. Chem.* 269:2365-2368; Kennedy, B.P., et al. (1995) *J. Biol. Chem.* 270: 22378-22385; Komada, M., et al. (1990) *Biochem. Biophys. Res. Commun.* 168: 1059-1065; Cupillard, L. et al. (1997) *J. Biol. Chem.* 272:15745-15752; and Nalefski,
30 E.A. et al. (1994) *J. Biol. Chem.* 269:18239-18249.)

Rat platelets secrete two types of phospholipases when stimulated, type II phospholipase A2 and phosphatidylserine-phospholipase A1 (PS-PLA1). PS-PLA1, also referred to as serine-phospholipid-selective phospholipase A (PLA), was purified from rat platelets and found to be a 55-kDa protein with active serine residues. The predicted 456-amino acid sequence contains a putative
35 short N-terminal signal sequence and a GX SXG motif (SEQ ID NO:10). PS-PLA1 shares about 30%

homology with mammalian lipases such as lipoprotein lipase, hepatic lipase, and pancreatic lipase. The residues that surround the active serine in PS-PLA1 include histidine and aspartic acid. The serine, histidine, and aspartic acid residues form a triad that is highly conserved among mammalian lipases. PS-PLA1 hydrolyzes a fatty acyl residue at the sn-1 position of lysophosphatidylserine and phosphatidylserine, and appears to be the first phospholipase that exclusively hydrolyzes the sn-1 position. (Sato, T. et al. (1997) J. Biol. Chem. 272:2192-2198.)

PLAs are implicated in a variety of disease processes. For example, PLAs are found in the pancreas, in cardiac tissue, and in inflammation-associated tissues. Pancreatic PLAs function in the digestion of dietary lipids and have been proposed to play a role in cell proliferation, smooth muscle contraction, and acute lung injury. Inflammatory PLAs are potent mediators of inflammatory processes and are highly expressed in serum and synovial fluids of patients with inflammatory disorders. Additionally, inflammatory PLAs are found in most human cell types and are expressed in diverse pathological processes such as septic shock, intestinal cancers, rheumatoid arthritis, and epidermal hyperplasia.

The discovery of new human phospholipases and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cancer, autoimmune/inflammatory disorders, and reproductive disorders.

SUMMARY OF THE INVENTION

The invention features substantially purified polypeptides, human phospholipases, referred to collectively as "HPPL" and individually as "HPPL-1," "HPPL-2," and "HPPL-3." In one aspect, the invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-3 and fragments thereof. The invention also includes a polypeptide comprising an amino acid sequence that differs by one or more conservative amino acid substitutions from an amino acid sequence selected from the group consisting of SEQ ID NO:1-3.

The invention further provides a substantially purified variant having at least 90% amino acid identity to at least one of the amino acid sequences selected from the group consisting of SEQ ID NO:1-3 and fragments thereof. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-3 and fragments thereof. The invention also includes an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-3 and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid

sequence selected from the group consisting of SEQ ID NO:1-3 and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1-3 and fragments thereof.

5 The invention also provides a method for detecting a polynucleotide in a sample containing nucleic acids, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence to at least one of the polynucleotides of the sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide in the sample. In one aspect, 10 the method further comprises amplifying the polynucleotide prior to hybridization.

 The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:4-6 and fragments thereof. The invention further provides an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide sequence selected from the group consisting of 15 SEQ ID NO:4-6 and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:4-6 and fragments thereof.

 The invention further provides an expression vector containing at least a fragment of the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-3. In another aspect, the expression vector is contained within a host cell. 20

 The invention also provides a method for producing a polypeptide, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing a polynucleotide of the invention under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

25 The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-3 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

 The invention further includes a purified antibody which binds to a polypeptide selected from the group consisting of SEQ ID NO:1-3 and fragments thereof. The invention also provides a purified 30 agonist and a purified antagonist to the polypeptide.

 The invention also provides a method for treating or preventing a disorder associated with decreased expression or activity of HPPL, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID 35 NO:1-3 and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention also provides a method for treating or preventing a disorder associated with increased expression or activity of HPPL, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-3 and fragments thereof.

BRIEF DESCRIPTION OF THE FIGURES AND TABLES

Figure 1 shows the amino acid sequence alignment between HPPL-1 (Incyte Clone number 2641779; SEQ ID NO:1) and group II phospholipase A-2 (GI 204319; SEQ ID NO:7), produced using the multisequence alignment program of LASERGENE software (DNASTAR, Madison WI).

Figures 2A, 2B, and 2C show the amino acid sequence alignment between HPPL-2 (Incyte Clone number 1430683; SEQ ID NO:2) and cytosolic phospholipase A2 (GI 508625; SEQ ID NO:8).

Figures 3A and 3B show the amino acid sequence alignment between HPPL-3 (Incyte Clone number 1316804; SEQ ID NO:3) and rat phosphatidylserine-phospholipase A1 (PS-PLA1) (GI 1817556; SEQ ID NO:9).

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding HPPL.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of HPPL.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding HPPL were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze HPPL, along with applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"HPPL" refers to the amino acid sequences of substantially purified HPPL obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of HPPL. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of HPPL either by directly interacting with HPPL or by acting on components of the biological pathway in which HPPL participates.

An "allelic variant" is an alternative form of the gene encoding HPPL. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding HPPL include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as HPPL or a polypeptide with at least one functional characteristic of HPPL. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding HPPL, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding HPPL. The encoded protein may also be "altered," and may contain deletions, insertions, or

substitutions of amino acid residues which produce a silent change and result in a functionally equivalent HPPL. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of HPPL is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of HPPL. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of HPPL either by directly interacting with HPPL or by acting on components of the biological pathway in which HPPL participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind HPPL polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures

n the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition containing a nucleic acid sequence which is complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic HPPL, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding HPPL or fragments of HPPL may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more Incyte Clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that, when made, least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
10	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
15	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
20	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
25	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "fragment" is a unique portion of HPPL or the polynucleotide encoding HPPL which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up

to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

A fragment of SEQ ID NO:4-6 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:4-6, for example, as distinct from any other sequence in the same genome. A fragment of SEQ ID NO:4-6 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:4-6 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:4-6 and the region of SEQ ID NO:4-6 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-3 is encoded by a fragment of SEQ ID NO:4-6. A fragment of SEQ ID NO:1-3 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-3. For example, a fragment of SEQ ID NO:1-3 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-3. The precise length of a fragment of SEQ ID NO:1-3 and the region of SEQ ID NO:1-3 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity

(e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example,
5 as defined by a particular SEQ ID number, or may be measured over a shorter length, for example,
over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at
least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous
nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported
by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a
10 length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode
similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes
in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid
sequences that all encode substantially the same protein.

15 The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to
the percentage of residue matches between at least two polypeptide sequences aligned using a
standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some
alignment methods take into account conservative amino acid substitutions. Such conservative
substitutions, explained in more detail above, generally preserve the hydrophobicity and acidity at the
20 site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default
parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e
sequence alignment program (described and referenced above). For pairwise alignments of
polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap
25 penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default
residue weight table. As with polynucleotide alignments, the percent identity is reported by
CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise
comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.9
30 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

35 *Word Size: 3*

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

“Human artificial chromosomes” (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term “humanized antibody” refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the “washing” step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention

include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C₀t or R₀t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate" refers to a change in the activity of HPPL. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of HPPL.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding

sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

“Peptide nucleic acid” (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

“Probe” refers to nucleic acid sequences encoding HPPL, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

“Primers” are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase

sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, *supra*. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding HPPL, or fragments thereof, or HPPL itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding

molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene

between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

5 A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence
10 identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human phospholipases (HPPL), the polynucleotides encoding HPPL, and the use of these compositions for the diagnosis, treatment, or
15 prevention of cancer, autoimmune/inflammatory disorders, and reproductive disorders.

Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding HPPL. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each HPPL were identified, and column 4 shows the cDNA libraries from
20 which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The Incyte clones in column 5 were used to assemble the consensus nucleotide sequence of each HPPL and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention:
25 column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The
30 methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

HPPL-1 has chemical and structural similarity with group II phospholipase A-2 from rat (GI 204319; SEQ ID NO:7). In particular, HPPL-1 and group II phospholipase A-2 share 46% identity. HPPL-2 has chemical and structural similarity with cytosolic phospholipase A2 from chicken (GI
35 508625; SEQ ID NO:8). In particular, HPPL-2 and cytosolic phospholipase A2 share 20% identity.

HPPL-3 has chemical and structural similarity with PS-PLA1 (GI 1817556; SEQ ID NO:9). In particular, HPPL-3 and PS-PLA1 share 80% identity. HPPL-3 and PS-PLA1 each contain a short N-terminal signal sequence, share the conserved potential lipase serine active site at residue I160 and the GX SXG (SEQ ID NO:10) consensus pattern.

5 The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding HPPL. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:4-6 and to distinguish between SEQ ID NO:4-6 and related polynucleotide sequences. The polypeptides
10 encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express HPPL as a fraction of total tissues expressing HPPL. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing HPPL as a fraction of total tissues expressing HPPL. Column 5 lists the vectors used to subclone each cDNA library. Of particular note is the expression of HPPL-1 (SEQ ID NO:1) in prostate, breast, and testicular tumors,
15 the expression of HPPL-2 (SEQ ID NO:2) in tumors of the ovary, breast, and brain, and the expression of HPPL-3 (SEQ ID NO:3) in gastrointestinal and reproductive tissue.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding HPPL were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows
20 the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

The invention also encompasses HPPL variants. A preferred HPPL variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the HPPL amino acid sequence, and which contains at least one functional or structural characteristic of HPPL.

25 The invention also encompasses polynucleotides which encode HPPL. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:4-6, which encodes HPPL.

The invention also encompasses a variant of a polynucleotide sequence encoding HPPL. In particular, such a variant polynucleotide sequence will have at least about 80%, or alternatively at
30 least about 90%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding HPPL. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:4-6 which has at least about 80%, or alternatively at least about 90%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of
35 SEQ ID NO:4-6. Any one of the polynucleotide variants described above can encode an amino acid

sequence which contains at least one functional or structural characteristic of HPPL.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding HPPL, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring HPPL, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode HPPL and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring HPPL under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding HPPL or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding HPPL and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode HPPL and HPPL derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding HPPL or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:4-6 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE

amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Perkin-Elmer). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding HPPL may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.)

Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060).

Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode HPPL may be cloned in recombinant DNA molecules that direct expression of HPPL, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express HPPL.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter HPPL-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding HPPL may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.)

Alternatively, HPPL itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of HPPL, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active HPPL, the nucleotide sequences encoding HPPL or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding HPPL. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding HPPL. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding HPPL and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding HPPL and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding HPPL. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding HPPL. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding HPPL can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding HPPL into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of

transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of HPPL are needed, e.g. for the production of antibodies, vectors which direct high level expression of HPPL may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of HPPL. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of HPPL. Transcription of sequences encoding HPPL may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding HPPL may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses HPPL in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of HPPL in cell lines is preferred. For example, sequences encoding HPPL can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *ap^r* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding HPPL is inserted within a marker gene sequence, transformed cells containing sequences encoding HPPL can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding HPPL under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding HPPL and that express HPPL may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or

chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of HPPL using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and
5 fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on HPPL is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and
10 Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding HPPL
15 include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding HPPL, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety
20 of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding HPPL may be cultured under
25 conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode HPPL may be designed to contain signal sequences which direct secretion of HPPL through a prokaryotic or eukaryotic cell membrane.

30 In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity.

35 Different host cells which have specific cellular machinery and characteristic mechanisms for

post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding HPPL may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric HPPL protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of HPPL activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the HPPL encoding sequence and the heterologous protein sequence, so that HPPL may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled HPPL may be achieved *in vitro* using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

Fragments of HPPL may be produced not only by recombinant means, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, *supra*, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of HPPL may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of HPPL and human phospholipases. In addition, the expression of HPPL is closely associated with cancerous, inflammation and the immune response, and with gastrointestinal and

reproductive tissues. Therefore, HPPL appears to play a role in cancer, autoimmune/inflammatory disorders, and reproductive disorders. In the treatment of disorders associated with increased HPPL expression or activity, it is desirable to decrease the expression or activity of HPPL. In the treatment of disorders associated with decreased HPPL expression or activity, it is desirable to increase the

5 expression or activity of HPPL.

Therefore, in one embodiment, HPPL or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HPPL.

Examples of such disorders include, but are not limited to, a cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of
10 the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis,
15 autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable
20 bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal,
25 parasitic, protozoal, and helminthic infections, and trauma; and a reproductive disorder, such as disorders of prolactin production; infertility, including tubal disease, ovulatory defects, and endometriosis; disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, endometrial and ovarian tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the breast, fibrocystic breast
30 disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia.

In another embodiment, a vector capable of expressing HPPL or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased
35 expression or activity of HPPL including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified HPPL in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HPPL including, but not limited to, those provided above.

5 In still another embodiment, an agonist which modulates the activity of HPPL may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of HPPL including, but not limited to, those listed above.

In a further embodiment, an antagonist of HPPL may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of HPPL. Examples of such
10 disorders include, but are not limited to, those cancer, autoimmune/inflammatory disorders, and reproductive disorders described above. In one aspect, an antibody which specifically binds HPPL may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express HPPL.

In an additional embodiment, a vector expressing the complement of the polynucleotide
15 encoding HPPL may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of HPPL including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made
20 by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of HPPL may be produced using methods which are generally known in the art.
25 In particular, purified HPPL may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind HPPL. Antibodies to HPPL may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit
30 dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with HPPL or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral
35 gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols,

polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to HPPL have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of HPPL amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to HPPL may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce HPPL-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for HPPL may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired

specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between HPPL and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering HPPL epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for HPPL. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of HPPL-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple HPPL epitopes, represents the average affinity, or avidity, of the antibodies for HPPL. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular HPPL epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the HPPL-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of HPPL, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington, DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of HPPL-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding HPPL, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding HPPL may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding HPPL. Thus, complementary molecules or fragments may be used to modulate HPPL activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from

various locations along the coding or control regions of sequences encoding HPPL.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding HPPL. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

Genes encoding HPPL can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding HPPL. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding HPPL. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding HPPL.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary

ligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis.

5 Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding HPPL. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

10 RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine,
15 and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient.

20 Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and
25 monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of HPPL, antibodies to HPPL, and mimetics, agonists, antagonists, or inhibitors of HPPL. The compositions
30 may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any
35 number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial,

intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration.

Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or

dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of HPPL, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example HPPL or fragments thereof, antibodies of HPPL, and agonists, antagonists or inhibitors of HPPL, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions

which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μ g to 100,000 μ g, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind HPPL may be used for the diagnosis of disorders characterized by expression of HPPL, or in assays to monitor patients being treated with HPPL or agonists, antagonists, or inhibitors of HPPL. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for HPPL include methods which utilize the antibody and a label to detect HPPL in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring HPPL, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of HPPL expression. Normal or standard values for HPPL expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to HPPL under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of HPPL expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation

between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding HPPL may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of HPPL may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of HPPL, and to monitor regulation of HPPL levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding HPPL or closely related molecules may be used to identify nucleic acid sequences which encode HPPL. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding HPPL, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the HPPL encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:4-6 or from genomic sequences including promoters, enhancers, and introns of the HPPL gene.

Means for producing specific hybridization probes for DNAs encoding HPPL include the cloning of polynucleotide sequences encoding HPPL or HPPL derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding HPPL may be used for the diagnosis of disorders associated with expression of HPPL. Examples of such disorders include, but are not limited to, a cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder, such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact

dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and a reproductive disorder, such as disorders of prolactin production; infertility, including tubal disease, ovulatory defects, and endometriosis; disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, endometrial and ovarian tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia. The polynucleotide sequences encoding HPPL may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered HPPL expression.

Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding HPPL may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding HPPL may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding HPPL in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of HPPL, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding HPPL, under conditions suitable for hybridization or amplification.

Standard hybridization may be quantified by comparing the values obtained from normal subjects

with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

5 Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

10 With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development
15 or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding HPPL may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding HPPL, or a fragment of a polynucleotide complementary to the polynucleotide encoding
20 HPPL, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

Methods which may also be used to quantify the expression of HPPL include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from
25 standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

30 In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and
35 monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding HPPL may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding HPPL on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, HPPL, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a

solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between HPPL and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with HPPL, or fragments thereof, and washed. Bound HPPL is then detected by methods well known in the art. Purified HPPL can also be coated directly onto plates for use in the aforementioned drug screening techniques.

Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding HPPL specifically compete with a test compound for binding HPPL. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with HPPL.

In additional embodiments, the nucleotide sequences which encode HPPL may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. [Attorney Docket No. PF-0625 P, filed October 27, 1998] and U.S. Ser. No. [Attorney Docket No. PF-0663 P, filed January 21, 1999], are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized

and lysed in phenol or in a suitable mixture of d naturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from th lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

5 Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA
10 purification kit (Ambion, Austin TX).

An alternative method of library construction was as follows:

The frozen tissue was homogenized and lysed in guanidinium isothiocyanate solution using a Brinkmann Homogenizer Polytron-PT 3000 (Brinkmann Instruments, Westbury NY). RNA was isolated as per Stratagene's RNA isolation protocol (Stratagene, La Jolla CA). RNA was extracted
15 twice with acid phenol, precipitated with sodium acetate and ethanol, resuspended in RNase-free water, and treated with DNase. Poly (A+) RNA was isolated using the OLIGOTEX kit (QIAGEN Inc, Chatsworth CA).

Poly(A+) RNA was used for cDNA synthesis and construction of the cDNA library according to the recommended protocols in the SUPERScript plasmid system (Life Technologies). The
20 cDNAs were fractionated on a SEPHAROSE CL4B column (Amersham Pharmacia Biotech), and those cDNAs exceeding 400 bp were ligated into pINCY (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into DH5 α competent cells or ELECTROMAX cells (Life Technologies).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA
25 libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, *supra*, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the
30 appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUEScript plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), or pINCY (Incyte
35 Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli*

cells including XL1-Blue, XL1-Blu MRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids were recovered from host cells by in vivo excision using the UNIZAP vector system
5 (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or
10 without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically
15 using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200
20 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides
25 were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques
30 disclosed in Example V.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools,
35 programs, and algorithms used, the second column provides brief descriptions thereof, the third

column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:4-6. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of

the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

100

5 The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

10 The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding HPPL occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma,
15 cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of HPPL Encoding Polynucleotides

20 The full length nucleic acid sequences of SEQ ID NO:4-6 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30
25 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

30 High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer
35 pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C,

2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

5 The concentration of DNA in each well was determined by dispensing 100 µl PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 µl of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the
10 concentration of DNA. A 5 µl to 10 µl aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

 The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and
15 sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site
20 overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

 The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following
25 parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing
30 primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

 In like manner, the nucleotide sequences of SEQ ID NO:4-6 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

35

VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:4-6 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments.

- 5 Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - 32 P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷
- 10 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

- The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16
- 15 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

VII. Microarrays

- 20 A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, supra.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After
- 25 hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

- Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise
- 30 the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking
- 35 followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al.

(1995) Science 270:467-470; Shalon, D. et al. (1996) Genom Res. 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the HPPL-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring HPPL. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of HPPL. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the HPPL-encoding transcript.

IX. Expression of HPPL

Expression and purification of HPPL is achieved using bacterial or virus-based expression systems. For expression of HPPL in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3).

Antibiotic resistant bacteria express HPPL upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of HPPL in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding HPPL by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, HPPL is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham

Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from HPPL at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified HPPL obtained by these methods can be used directly in the following activity assay.

X. Demonstration of HPPL Activity

HPPL activity can be demonstrated by an in vitro hydrolysis assay with vesicles containing 1-palmitoyl-2-[1-¹⁴C]oleoyl phosphatidylcholine (Sigma-Aldrich). HPPL phospholipase A₂ activity is demonstrated by analysis of the cleavage products isolated from the hydrolysis reaction mixture.

Vesicles containing 1-palmitoyl-2-[1-¹⁴C]oleoyl phosphatidylcholine (Amersham Pharmacia Biotech.) are prepared by mixing 2.0 µCi of the radiolabeled phospholipid with 12.5 mg of unlabeled 1-palmitoyl-2-oleoyl phosphatidylcholine and drying the mixture under N₂. 2.5 ml of 150 mM

Tris-HCl, pH 7.5, is added, and the mixture is sonicated and centrifuged. The supernatant may be stored at 4 °C. The final reaction mixtures contain 0.25 ml of Hanks buffered salt solution supplemented with 2.0 mM taurochenodeoxycholate, 1.0% bovine serum albumin, 1.0 mM CaCl₂, pH 7.4, 150 µg of 1-palmitoyl-2-[1-¹⁴C]oleoyl phosphatidylcholine vesicles, and various amount of HPPL diluted in PBS. After incubation for 30 min at 37 °C, 20 µg each of lyso-phosphatidylcholine and oleic acid are added as carriers and each sample is extracted for total lipids. The lipids are separated by thin layer chromatography using a two solvent system of chloroform:methanol:acetic acid:water (65:35:8:4) until the solvent front is halfway up the plate. The process is then continued with hexane:ether:acetic acid (86:16:1) until the solvent front is at the top of the plate. The lipid-containing areas are visualized with I₂ vapor; the spots are scraped, and their radioactivity is determined by scintillation counting. The amount of radioactivity released as fatty acids will increase as a greater amount of HPPL is added to the assay mixture while the amount of radioactivity released as lyso-phosphatidylcholine will remain low. This demonstrates that HPPL cleaves at the *sn*-2 and not the *sn*-1 position, as characteristic of phospholipase A₂ activity.

Alternatively, HPPL activity is measured by the hydrolysis of a fatty acyl residue at the *sn*-1 position of phosphatidylserine. HPPL is combined with the Tritium [³H] labeled substrate phosphatidylserine at stoichiometric quantities in a suitable buffer. Following an appropriate incubation time, the hydrolyzed reaction products are separated from the substrates by chromatographic methods. The amount of acylglycerophosphoserine produced is measured by counting tritiated product with the help of a scintillation counter. Various control groups are set up to account for background noise and unincorporated substrate. The final counts represent the tritiated

enzym product [3H]-acylgly rophosphos rine, which is directly proportional to the activity of HPPL in biological samples.

XI. Functional Assays

HPPL function is assessed by expressing the sequences encoding HPPL at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT (Life Technologies) and pCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of HPPL on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding HPPL and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding HPPL and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of HPPL Specific Antibodies

HPPL substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the HPPL amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-HPPL activity by, for example, binding the peptide or HPPL to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIII. Purification of Naturally Occurring HPPL Using Specific Antibodies

Naturally occurring or recombinant HPPL is substantially purified by immunoaffinity chromatography using antibodies specific for HPPL. An immunoaffinity column is constructed by covalently coupling anti-HPPL antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing HPPL are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of HPPL (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/HPPL binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and HPPL is collected.

XIV. Identification of Molecules Which Interact with HPPL

HPPL, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled HPPL, washed, and any wells with labeled HPPL complex are assayed. Data obtained using different concentrations of HPPL are used to calculate values for the number, affinity, and association of HPPL with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention

will be apparent to those skilled in the art without departing from the scope and spirit of the invention.

Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments.

Indeed, various modifications of the described modes for carrying out the invention which are

- 5 obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Polypeptide SEQ ID NO:	Polynucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	4	2641779	LUNGTUT08	2641779H1 (LUNGTUT08), 3980344H1 (LUNGTUT08), 2203422F6 (SPLNFET02), 701224H1 (SYNORAT03)
2	5	1430683	SINTBST01	1430683H1 (SINTBST01), 1430683F6 (SINTBST01), 4591542H1 (MASTTX01), 1654109X12 (PROSTUT08), 836234R6 (PROSNOT07), 1511418F6 (LUNGNOT14), 1513169F6 (PANCUT01)
3	6	1316804	BLADTUT02	1316804H1 (BLADTUT02), 1316804T6 (BLADTUT02), 2965020H1 (SCORNOT04), 3539795H1 (SEMVN04), SBFA00393F1, SBFA01391F1, SBFA03702F1, SBFA04815F1, SBFA04928F1

Table 2

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential glycosylation sites	Signature Sequences, Motifs and Domains	Homologous Sequences	Analytical Methods & Databases
1	145	S102 T33 S98	N89	Signal peptide: M1-I16; M1-G21 Active site histidine region: Y44-Q92 Active site aspartic acid region: Y88-G140 phospholipase A2 active site signature: G21-C145	Group IId secreted phospholipase A2, group II (g5359708) <u>Mus musculus</u> Group II Phospholipase A-2 (g204319) <u>Rattus norvegicus</u>	PROFILERSCAN HMMER-PFAM HMMER BLIMPS-BLOCKS BLIMPS-PRINTS SPSCAN BLAST-GenBank
2	605	S72 S78 S186 T282 S293 T430 T440 T506 T518 T540 S552 S566 S104 T397 T426	N456 N550		Cytosolic phospholipase A2 beta (g3811347) <u>Homo sapiens</u> Cytosolic phospholipase A2 (g508625) <u>Gallus gallus</u>	BLAST-GenBank

Table 2 cont.

Poly-peptide SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential glycosylation sites	Signature Sequences, Motifs and Domains	Homologous Sequences	Analytical Methods & Databases
3	456	S23 S99 T187 S284 S369 S404	N79 N365	Signal peptide: M1-G25; M1-A27 Lipase serine active site: I160 Triglyceride lipase GXSG (SEQ ID NO:10) consensus pattern: G164-G168 Putative lipase signature: T47-C245; C258-C296 Triacylglycerol lipase family signature: D72-F91; I117-A132; S159-Q177; C258- N273	Serine phospholipid- specific phospholipase A (g1817556) <u>Rattus norvegicus</u>	HMME-PFAM BLIMPS-PRINTS HMMR SPSCAN BLAST-GenBank MOTIFS

Table 3

Nucleotide SEQ ID NO:	Selected Fragment (s)	Tissue Expression	Disease or Condition (Fraction of Total)	Vector
4	14-73	Reproductive Cardiovascular	Cancer and proliferating tissue (0.890) Inflammation/Trauma (0.220)	pINCY
5	95-154	Reproductive Nervous	Cancer and proliferating tissue (0.720) Immune Response (0.240)	PBLUESCRIPT
6	1148-1192	Reproductive Gastrointestinal	Cancer (0.500) Inflammation and Immune Response (0.420)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Description
1	LUNGUTUT08	Library was constructed using RNA isolated from lung tumor tissue removed from a 63-year old Caucasian male during a right upper lobectomy with fiberoptic bronchoscopy. Pathology indicated a grade 3 adenocarcinoma. Patent history included rectal cancer and tobacco abuse. Family history included stomach cancer and lung cancer.
2	SINTBST01	Library was constructed using RNA isolated from ileum tissue obtained from an 18-year old Caucasian female removed during bowel anastomosis. Pathology indicated Crohn's disease of the ileum, involving 15 cm of the small bowel.
3	BLADTUT02	Library was constructed using RNA isolated from bladder tumor tissue removed from an 80-year-old Caucasian female during a radical cystectomy and lymph node excision. Pathology indicated grade 3 invasive transitional cell carcinoma. Family history included acute renal failure, osteoarthritis, and atherosclerosis.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises as least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value= 1.0E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value= 1.0E-8 or less Full Length sequences: fastx score= 100 or greater
BLIMPS	A BLOCKS IMPROVED Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score= 1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score= 10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality score \geq GCG-specified "HIGH" value for that particular Prosite motif. Generally, score = 1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score = 120 or greater; Match length = 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score = 3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <u>supra</u> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. A substantially purified polypeptid comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1 and SEQ ID NO:3 and fragm nts thereof.

2. A substantially purified variant having at least 90% amino acid sequence identity to the amino acid sequence of claim 1.

3. An isolated and purified polynucleotide encoding the polypeptide of claim 1.

4. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 3.

5. An isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide of claim 3.

6. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 3.

7. A method for detecting a polynucleotide, the method comprising the steps of:

(a) hybridizing the polynucleotide of claim 6 to at least one nucleic acid in a sample, thereby forming a hybridization complex; and

(b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of the polynucleotide in the sample.

8. The method of claim 7 further comprising amplifying the polynucleotide prior to hybridization.

9. An isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:4 and SEQ ID NO:5 and fragments thereof.

10. An isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide of claim 9.

11. An isolated and purified polynucleotide having a sequence which is complementary

to the polynucleotide of claim 9.

12. An expression vector comprising at least a fragment of the polynucleotide of claim 3.

5 13. A host cell comprising the expression vector of claim 12.

14. A method for producing a polypeptide, the method comprising the steps of:

a) culturing the host cell of claim 13 under conditions suitable for the expression of the polypeptide; and

10 b) recovering the polypeptide from the host cell culture.

15. A pharmaceutical composition comprising the polypeptide of claim 1 in conjunction with a suitable pharmaceutical carrier.

15 16. A purified antibody which specifically binds to the polypeptide of claim 1.

17. A purified agonist of the polypeptide of claim 1.

18. A purified antagonist of the polypeptide of claim 1.

20

19. A method for treating or preventing a disorder associated with decreased expression or activity of HPPL, the method comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 15.

25 20. A method for treating or preventing a disorder associated with increased expression or activity of HPPL, the method comprising administering to a subject in need of such treatment an effective amount of the antagonist of claim 18.

1	M	E	L	A	L	L	C	G	L	V	V	M	A	-	G	V	I	P	I	Q	G	G	I	L	N	L	N	K	M	V	2641779
1	M	K	V	L	L	L	L	A	V	V	I	M	A	F	G	S	I	Q	V	Q	G	S	L	L	E	F	G	Q	M	I	GI 204319
30	K	Q	V	T	G	K	M	P	I	L	S	Y	W	P	Y	G	C	H	C	G	L	G	G	R	G	Q	P	K	D	A	2641779
31	L	F	K	T	G	K	R	A	D	V	S	Y	G	F	Y	G	C	H	C	G	V	G	G	R	G	S	P	K	D	A	GI 204319
60	T	D	W	C	C	Q	T	H	D	C	C	Y	D	H	L	K	T	Q	G	C	G	I	Y	K	D	Y	Y	R	Y	N	2641779
61	T	D	W	C	C	V	T	H	D	C	C	Y	N	R	L	E	K	R	G	C	G	T	K	F	L	T	Y	K	F	S	GI 204319
90	F	S	Q	G	N	I	H	C	S	D	K	G	S	W	C	E	Q	Q	L	C	A	C	D	K	E	V	A	F	C	L	2641779
91	Y	R	G	G	Q	I	S	C	S	T	N	Q	D	S	C	R	K	Q	L	C	Q	C	D	K	A	A	A	E	C	F	GI 204319
120	K	R	N	L	D	T	Y	Q	K	R	L	R	F	Y	W	R	P	H	C	R	G	Q	T	P	G	C	2641779				
121	A	R	N	K	K	S	Y	S	L	K	Y	Q	F	Y	L	N	K	F	C	K	G	K	T	P	S	C	GI 204319				

FIGURE 1

1	- - - - - M I F V E - - L S P T L A L C L E R V A S H	1430683
11	M S F I D P Y Q H I V V E H Q Y S H V F T V T V - R K A T N	GI 508625
21	L T D T G L - - - - - L V L F C P A P - - C P - -	1430683
30	V T K G A I G D M L D T P D P Y V E L F I P S A P D C R K R	GI 508625
37	- - - - - - - - - - - F F F F F E M - - - - -	1430683
60	T K H F N N D V N P V W N E T F F E F I L D P N Q D N V L E V	GI 508625
44	- - - - - - - - - - E S L S V A Q A G V Q W R D L G S L Q P	1430683
90	T L M D A N Y V M D E T L G M A T F P I S S L K L G E K K E	GI 508625
64	P P L G F K R F S C L S L P S S W D Y R L R E L A V R L G F	1430683
120	V Q L T F N N V T E M T L E L S L E V C - S S T D L R F S M	GI 508625
94	G P C A E E Q A F L S R R K Q V V A A L R Q A L Q L D G -	1430683
149	A L C D E E K K F R Q Q R K D N I M Q S M K S F L G E E N S	GI 508625
123	- D L Q E D - E I P V V A I M A T G G G I R A M T S L Y G Q	1430683
179	K N L T T S R D V P V I A V L G S G G G F R A M V G F A G V	GI 508625
151	L A G L K E L G L L D C V S Y I T G A S G S T W A L A N L Y	1430683
209	M K A L Y E S G V L D C A T Y I A G L S G S T W Y M S T L Y	GI 508625
181	E D P E W S Q K D L A G P T E L L K T Q V T K N K L G V L A	1430683
239	S H P D F P E K G P K E I N Q E L M N S V S H N P L L L L T	GI 508625

FIGURE 2A

3/6

211	P	S	Q	L	Q	R	Y	R	Q	E	L	A	E	R	A	R	L	G	Y	P	S	C	F	T	N	L	W	A	-	L	1430683										
269	P	Q	K	V	K	R	Y	I	E	A	L	W	N	K	K	S	S	G	Q	P	V	T	F	T	D	I	F	G	M	L	GI 508625										
240	I	N	E	A	L	L	H	D	E	P	H	D	H	K	L	S	D	Q	R	E	A	L	S	H	G	Q	N	P	L	P	1430683										
299	I	G	E	T	L	I	H	N	-	R	M	D	T	T	L	S	D	M	K	E	K	V	S	E	A	Q	C	A	L	P	GI 508625										
270	I	Y	C	A	L	N	T	K	G	Q	S	L	T	T	F	E	F	G	E	W	C	E	F	S	P	Y	E	V	G	F	1430683										
328	L	F	T	C	L	H	V	K	P	D	-	V	S	E	L	M	F	A	D	W	V	E	F	S	P	Y	E	I	G	M	GI 508625										
300	P	K	Y	G	A	F	I	P	S	E	L	F	G	S	E	F	F	M	G	Q	L	M	K	R	L	P	E	S	R	I	1430683										
357	A	K	Y	G	T	F	M	S	P	D	L	F	G	S	K	F	F	M	G	T	V	V	K	K	Y	S	E	N	P	L	GI 508625										
330	C	F	L	E	G	I	W	S	N	L	Y	A	A	N	L	Q	D	S	L	-	-	-	-	-	-	-	-	-	-	Y	1430683										
387	H	F	L	M	G	V	W	G	S	A	F	S	I	L	F	N	R	V	L	G	V	S	N	S	Q	N	K	G	P	T	GI 508625										
350	W	A	S	E	P	S	Q	F	W	-	D	R	W	V	R	N	Q	A	N	L	D	K	E	Q	V	P	L	L	K	I	1430683										
417	M	E	E	E	L	E	N	I	R	L	K	H	L	V	S	N	D	S	S	D	S	E	D	E	S	Q	H	P	K	G	GI 508625										
379	E	E	P	P	S	T	A	-	-	-	-	-	-	-	-	-	-	-	-	G	R	I	A	E	F	F	T	D	L	T	W	R	1430683								
447	T	E	N	S	E	A	N	E	E	Y	Q	N	S	S	Q	E	S	W	V	Q	R	M	L	M	A	L	V	G	D	S	GI 508625										
400	P	L	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	A	Q	A	T	H	N	F	L	R	G	L	H	F	H	K	D	Y	F	Q	H	P	1430683
477	A	L	F	N	T	R	E	G	R	A	G	K	V	H	N	F	M	L	G	L	N	L	N	S	C	Y	P	L	S	P	GI 508625										
423	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1430683		
507	L	A	D	L	L	T	Q	E	S	V	E	E	D	E	L	D	A	A	V	A	D	P	D	E	F	E	R	I	Y	E	GI 508625										

FIGURE 2B

4/6

445 - - - - - H L C L L D V G Y L I N T S C L P L L Q P T R 1430683
537 P L D V K S K K I H I V D S G L T F N L P Y P L I L R P Q R GI 508625

468 D V D L I L S L D Y N L H G A - - - - - F Q Q L Q L L G R F 1430683
567 G V D L I I S F D F S A R P S D S S P P F K E I L L A E K W GI 508625

493 C Q E Q G I P F P P I S P S P E E Q L Q P R E C H T F S D P 1430683
597 A K M N K L P F P K I D P N V F D R E G L K E C Y V F K P K GI 508625

523 T C P G A - - - - - P A V L H F - - - - - S S G V R 1430683
627 D T S S E K D C P T I I H F V L A N I N F R K Y K A P G L P GI 508625

539 R T P E E A A A - G E V N L - S S S D S P Y H Y T K V T Y S 1430683
657 R E S K E E K D F A D F D I F D D P N T P F S T F N F Q Y P GI 508625

567 Q E D V D K L L H L T H Y N V C N N Q E Q L L E A L R Q A V 1430683
687 N E A F K R L H D L M E F N T L N N L D V I K Q A M M E S I GI 508625

597 Q R R R Q R R P H 1430683
717 E Y R K E N P S R C S V S L S S V E A R R F F N K N N L N N GI 508625

605
747 H T 1430683
GI 508625

FIGURE 2C

5/6

1	M	P	P	G	P	W	E	S	C	F	W	V	G	G	L	I	L	W	L	S	V	G	S	S	G	D	A	P	P	T	1316804
1	M	C	P	G	L	W	G	T	C	F	W	L	W	G	S	L	L	W	L	S	I	G	R	S	G	N	V	P	P	T	g1817556
31	P	Q	P	K	C	A	D	F	Q	S	A	N	L	F	E	G	T	D	L	K	V	Q	F	L	L	F	V	P	S	N	1316804
31	T	Q	P	K	C	T	D	F	Q	S	A	N	L	L	R	G	T	N	L	K	V	Q	F	L	L	F	T	P	S	D	g1817556
61	P	S	C	G	Q	L	V	E	G	S	S	D	L	Q	N	S	G	F	N	A	T	L	G	T	K	L	I	I	H	G	1316804
61	P	G	C	G	Q	L	V	E	D	S	D	I	R	N	S	E	F	N	A	S	L	G	T	K	L	I	I	H	G	g1817556	
91	F	R	V	L	G	T	K	P	S	W	I	D	T	F	I	R	T	L	L	R	A	T	N	A	N	V	I	A	V	D	1316804
91	F	R	A	L	G	T	K	P	S	W	I	N	K	F	I	R	A	L	L	R	A	A	D	A	N	V	I	A	V	D	g1817556
121	W	I	Y	G	S	T	G	V	Y	F	S	A	V	K	N	V	I	K	L	S	L	E	I	S	L	F	L	N	K	L	1316804
121	W	V	Y	G	S	T	G	M	Y	F	S	A	V	E	N	V	V	K	L	S	L	E	I	S	R	F	L	S	K	L	g1817556
151	L	V	L	G	V	S	E	S	S	I	H	I	I	G	V	S	L	G	A	H	V	G	G	M	V	G	Q	L	F	G	1316804
151	L	E	L	G	V	S	E	S	S	I	H	I	I	G	V	S	L	G	A	H	V	G	G	M	V	G	H	F	Y	K	g1817556
181	G	Q	L	G	Q	I	T	G	L	D	P	A	G	P	E	Y	T	R	A	S	V	E	E	R	L	D	A	G	D	A	1316804
181	G	Q	L	G	R	I	T	G	L	D	P	A	G	P	E	Y	T	R	A	S	L	E	E	R	L	D	S	G	D	A	g1817556
211	L	F	V	E	A	I	H	T	D	T	D	N	L	G	I	R	I	P	V	G	H	V	D	Y	F	V	N	G	G	Q	1316804
211	L	F	V	E	A	I	H	T	D	T	D	N	L	G	I	R	I	P	V	G	H	V	D	Y	F	V	N	G	G	Q	g1817556

FIGURE 3A

6/6

241	D	Q	P	G	C	P	T	F	F	Y	A	G	Y	S	Y	L	I	C	D	H	M	R	A	V	H	L	Y	I	S	A	1316804
241	D	Q	P	G	C	P	A	F	I	H	A	G	Y	S	Y	L	I	C	D	H	M	R	A	V	H	L	Y	I	S	A	g1817556
271	L	E	N	S	C	P	L	M	A	F	P	C	A	S	Y	K	A	F	L	A	G	R	C	L	D	C	F	N	P	F	1316804
271	L	E	N	T	C	P	L	M	A	F	P	C	A	S	Y	K	A	F	L	A	G	D	C	L	D	C	F	N	P	F	g1817556
301	L	L	S	C	P	R	I	G	L	V	E	Q	G	G	V	K	I	E	P	L	P	K	E	V	K	V	Y	L	L	T	1316804
301	L	L	S	C	P	R	I	G	L	V	E	R	G	G	V	K	I	E	P	L	P	K	E	V	R	V	Y	L	Q	T	g1817556
331	T	S	S	A	P	Y	C	M	H	S	L	V	E	F	H	L	K	E	L	R	N	K	D	T	N	I	E	V	T	1316804	
331	T	S	S	A	P	Y	C	V	H	S	L	V	E	F	N	L	K	E	K	R	K	D	T	S	I	E	V	T	g1817556		
361	F	L	S	S	N	I	T	S	S	S	K	I	T	I	P	K	Q	Q	R	Y	G	K	G	I	I	A	H	A	T	P	1316804
361	F	L	G	N	N	V	T	S	S	V	K	I	T	I	P	K	D	H	L	E	G	R	G	I	I	A	H	Q	N	P	g1817556
391	Q	C	Q	I	N	Q	V	K	F	K	F	Q	S	S	N	R	V	W	K	K	D	R	T	T	I	I	G	K	F	C	1316804
391	H	C	Q	I	N	Q	V	K	L	K	F	H	I	S	S	R	V	W	R	K	D	R	T	P	I	V	G	T	F	C	g1817556
421	T	A	L	L	P	V	N	D	R	E	K	M	V	C	L	P	E	P	V	N	L	Q	A	S	V	T	V	S	C	D	1316804
421	T	A	P	L	P	V	N	D	S	K	K	T	V	C	I	P	E	P	V	R	L	Q	V	S	M	A	V	L	R	D	g1817556
451	L	K	I	A	C	V	1316804																								
451	L	K	M	A	C	V	g1817556																								

FIGURE 3B

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.

HILLMAN, Jennifer L.

BANDMAN, Olga

GUEGLER, Karl J.

CORLEY, Neil C.

BAUGHN, Mariah R.

AZIMZAI, Yalda

LAL, Preeti

LY, Dyung Aina M.

<120> HUMAN PHOSPHOLIPASES

<130> PF-0625 PCT

<140> To Be Assigned

<141> Herewith

<150> 09/181,317; unassigned; 09/234,726; unassigned

<151> 1998-10-27; 1998-10-27; 1999-01-21; 1999-01-21

<160> 9

<170> PERL Program

<210> 1

<211> 145

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2641779CD1

<400> 1

Met	Glu	Leu	Ala	Leu	Leu	Cys	Gly	Leu	Val	Val	Met	Ala	Gly	Val
1				5					10					15
Ile	Pro	Ile	Gln	Gly	Gly	Ile	Leu	Asn	Leu	Asn	Lys	Met	Val	Lys
			20						25					30
Gln	Val	Thr	Gly	Lys	Met	Pro	Ile	Leu	Ser	Tyr	Trp	Pro	Tyr	Gly
			35						40					45
Cys	His	Cys	Gly	Leu	Gly	Gly	Arg	Gly	Gln	Pro	Lys	Asp	Ala	Thr
			50						55					60
Asp	Trp	Cys	Cys	Gln	Thr	His	Asp	Cys	Cys	Tyr	Asp	His	Leu	Lys
			65						70					75
Thr	Gln	Gly	Cys	Gly	Ile	Tyr	Lys	Asp	Tyr	Tyr	Arg	Tyr	Asn	Phe
			80						85					90
Ser	Gln	Gly	Asn	Ile	His	Cys	Ser	Asp	Lys	Gly	Ser	Trp	Cys	Glu
			95						100					105
Gln	Gln	Leu	Cys	Ala	Cys	Asp	Lys	Glu	Val	Ala	Phe	Cys	Leu	Lys
			110						115					120
Arg	Asn	Leu	Asp	Thr	Tyr	Gln	Lys	Arg	Leu	Arg	Phe	Tyr	Trp	Arg
			125						130					135

Pro His Cys Arg Gly Gln Thr Pro Gly Cys
140 145

<210> 2

<211> 605

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1430683CD1

<400> 2

Met	Ile	Phe	Val	Glu	Leu	Ser	Pro	Thr	Leu	Ala	Leu	Cys	Leu	Glu	
1				5					10					15	
Arg	Val	Ala	Ser	His	Leu	Thr	Asp	Thr	Gly	Leu	Leu	Val	Leu	Phe	
			20						25					30	
Cys	Pro	Ala	Pro	Cys	Pro	Phe	Phe	Phe	Phe	Phe	Glu	Met	Glu	Ser	
			35						40					45	
Leu	Ser	Val	Ala	Gln	Ala	Gly	Val	Gln	Trp	Arg	Asp	Leu	Gly	Ser	
			50						55					60	
Leu	Gln	Pro	Pro	Pro	Leu	Gly	Phe	Lys	Arg	Phe	Ser	Cys	Leu	Ser	
			65						70					75	
Leu	Pro	Ser	Ser	Trp	Asp	Tyr	Arg	Leu	Arg	Glu	Leu	Ala	Val	Arg	
			80						85					90	
Leu	Gly	Phe	Gly	Pro	Cys	Ala	Glu	Glu	Gln	Ala	Phe	Leu	Ser	Arg	
			95						100					105	
Arg	Lys	Gln	Val	Val	Ala	Ala	Ala	Leu	Arg	Gln	Ala	Leu	Gln	Leu	
			110						115					120	
Asp	Gly	Asp	Leu	Gln	Glu	Asp	Glu	Ile	Pro	Val	Val	Ala	Ile	Met	
			125						130					135	
Ala	Thr	Gly	Gly	Gly	Ile	Arg	Ala	Met	Thr	Ser	Leu	Tyr	Gly	Gln	
			140						145					150	
Leu	Ala	Gly	Leu	Lys	Glu	Leu	Gly	Leu	Leu	Asp	Cys	Val	Ser	Tyr	
			155						160					165	
Ile	Thr	Gly	Ala	Ser	Gly	Ser	Thr	Trp	Ala	Leu	Ala	Asn	Leu	Tyr	
			170						175					180	
Glu	Asp	Pro	Glu	Trp	Ser	Gln	Lys	Asp	Leu	Ala	Gly	Pro	Thr	Glu	
			185						190					195	
Leu	Leu	Lys	Thr	Gln	Val	Thr	Lys	Asn	Lys	Leu	Gly	Val	Leu	Ala	
			200						205					210	
Pro	Ser	Gln	Leu	Gln	Arg	Tyr	Arg	Gln	Glu	Leu	Ala	Glu	Arg	Ala	
			215						220					225	
Arg	Leu	Gly	Tyr	Pro	Ser	Cys	Phe	Thr	Asn	Leu	Trp	Ala	Leu	Ile	
			230						235					240	
Asn	Glu	Ala	Leu	Leu	His	Asp	Glu	Pro	His	Asp	His	Lys	Leu	Ser	
			245						250					255	
Asp	Gln	Arg	Glu	Ala	Leu	Ser	His	Gly	Gln	Asn	Pro	Leu	Pro	Ile	
			260						265					270	
Tyr	Cys	Ala	Leu	Asn	Thr	Lys	Gly	Gln	Ser	Leu	Thr	Thr	Phe	Glu	
			275						280					285	
Phe	Gly	Glu	Trp	Cys	Glu	Phe	Ser	Pro	Tyr	Glu	Val	Gly	Phe	Pro	
			290						295					300	

Lys Tyr Gly Ala Phe Ile Pro Ser Glu Leu Phe Gly Ser Glu Phe	305	310	315
Phe Met Gly Gln Leu Met Lys Arg Leu Pro Glu Ser Arg Ile Cys	320	325	330
Phe Leu Glu Gly Ile Trp Ser Asn Leu Tyr Ala Ala Asn Leu Gln	335	340	345
Asp Ser Leu Tyr Trp Ala Ser Glu Pro Ser Gln Phe Trp Asp Arg	350	355	360
Trp Val Arg Asn Gln Ala Asn Leu Asp Lys Glu Gln Val Pro Leu	365	370	375
Leu Lys Ile Glu Glu Pro Pro Ser Thr Ala Gly Arg Ile Ala Glu	380	385	390
Phe Phe Thr Asp Leu Leu Thr Trp Arg Pro Leu Ala Gln Ala Thr	395	400	405
His Asn Phe Leu Arg Gly Leu His Phe His Lys Asp Tyr Phe Gln	410	415	420
His Pro His Phe Ser Thr Trp Lys Ala Thr Thr Leu Asp Gly Leu	425	430	435
Pro Asn Gln Leu Thr Pro Ser Glu Pro His Leu Cys Leu Leu Asp	440	445	450
Val Gly Tyr Leu Ile Asn Thr Ser Cys Leu Pro Leu Leu Gln Pro	455	460	465
Thr Arg Asp Val Asp Leu Ile Leu Ser Leu Asp Tyr Asn Leu His	470	475	480
Gly Ala Phe Gln Gln Leu Gln Leu Leu Gly Arg Phe Cys Gln Glu	485	490	495
Gln Gly Ile Pro Phe Pro Pro Ile Ser Pro Ser Pro Glu Glu Gln	500	505	510
Leu Gln Pro Arg Glu Cys His Thr Phe Ser Asp Pro Thr Cys Pro	515	520	525
Gly Ala Pro Ala Val Leu His Phe Ser Ser Gly Val Arg Arg Thr	530	535	540
Pro Glu Glu Ala Ala Ala Gly Glu Val Asn Leu Ser Ser Ser Asp	545	550	555
Ser Pro Tyr His Tyr Thr Lys Val Thr Tyr Ser Gln Glu Asp Val	560	565	570
Asp Lys Leu Leu His Leu Thr His Tyr Asn Val Cys Asn Asn Gln	575	580	585
Glu Gln Leu Leu Glu Ala Leu Arg Gln Ala Val Gln Arg Arg Arg	590	595	600
Gln Arg Arg Pro His			605

<210> 3

<211> 456

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1316804CD1

<400> 3

```

Met Pro Pro Gly Pro Trp Glu Ser Cys Phe Trp Val Gly Gly Leu
 1          5          10          15
Ile Leu Trp Leu Ser Val Gly Ser Ser Gly Asp Ala Pro Pro Thr
          20          25          30
Pro Gln Pro Lys Cys Ala Asp Phe Gln Ser Ala Asn Leu Phe Glu
          35          40          45
Gly Thr Asp Leu Lys Val Gln Phe Leu Leu Phe Val Pro Ser Asn
          50          55          60
Pro Ser Cys Gly Gln Leu Val Glu Gly Ser Ser Asp Leu Gln Asn
          65          70          75
Ser Gly Phe Asn Ala Thr Leu Gly Thr Lys Leu Ile Ile His Gly
          80          85          90
Phe Arg Val Leu Gly Thr Lys Pro Ser Trp Ile Asp Thr Phe Ile
          95          100          105
Arg Thr Leu Leu Arg Ala Thr Asn Ala Asn Val Ile Ala Val Asp
          110          115          120
Trp Ile Tyr Gly Ser Thr Gly Val Tyr Phe Ser Ala Val Lys Asn
          125          130          135
Val Ile Lys Leu Ser Leu Glu Ile Ser Leu Phe Leu Asn Lys Leu
          140          145          150
Leu Val Leu Gly Val Ser Glu Ser Ser Ile His Ile Ile Gly Val
          155          160          165
Ser Leu Gly Ala His Val Gly Gly Met Val Gly Gln Leu Phe Gly
          170          175          180
Gly Gln Leu Gly Gln Ile Thr Gly Leu Asp Pro Ala Gly Pro Glu
          185          190          195
Tyr Thr Arg Ala Ser Val Glu Glu Arg Leu Asp Ala Gly Asp Ala
          200          205          210
Leu Phe Val Glu Ala Ile His Thr Asp Thr Asp Asn Leu Gly Ile
          215          220          225
Arg Ile Pro Val Gly His Val Asp Tyr Phe Val Asn Gly Gly Gln
          230          235          240
Asp Gln Pro Gly Cys Pro Thr Phe Phe Tyr Ala Gly Tyr Ser Tyr
          245          250          255
Leu Ile Cys Asp His Met Arg Ala Val His Leu Tyr Ile Ser Ala
          260          265          270
Leu Glu Asn Ser Cys Pro Leu Met Ala Phe Pro Cys Ala Ser Tyr
          275          280          285
Lys Ala Phe Leu Ala Gly Arg Cys Leu Asp Cys Phe Asn Pro Phe
          290          295          300
Leu Leu Ser Cys Pro Arg Ile Gly Leu Val Glu Gln Gly Gly Val
          305          310          315
Lys Ile Glu Pro Leu Pro Lys Glu Val Lys Val Tyr Leu Leu Thr
          320          325          330
Thr Ser Ser Ala Pro Tyr Cys Met His His Ser Leu Val Glu Phe
          335          340          345
His Leu Lys Glu Leu Arg Asn Lys Asp Thr Asn Ile Glu Val Thr
          350          355          360
Phe Leu Ser Ser Asn Ile Thr Ser Ser Ser Lys Ile Thr Ile Pro
          365          370          375
Lys Gln Gln Arg Tyr Gly Lys Gly Ile Ile Ala His Ala Thr Pro
          380          385          390
Gln Cys Gln Ile Asn Gln Val Lys Phe Lys Phe Gln Ser Ser Asn
          395          400          405

```

Arg Val Trp Lys Lys Asp Arg Thr Thr Ile Ile Gly Lys Phe Cys
 410 415 420
 Thr Ala Leu Leu Pro Val Asn Asp Arg Glu Lys Met Val Cys Leu
 425 430 435
 Pro Glu Pro Val Asn Leu Gln Ala Ser Val Thr Val Ser Cys Asp
 440 445 450
 Leu Lys Ile Ala Cys Val
 455

<210> 4

<211> 592

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 2641779CB1

<400> 4

tctgcctcca ctgctctgtg ctgggatcat ggaacttgca ctgctgtgtg ggctgggtgt 60
 gatggctggt gtgattccaa tccagggcgg gatcctgaac ctgaacaaga tggtaagca 120
 agtgactggg aaaatgccca tcctctccta ctggccctac ggctgtcact gcggactagg 180
 tggcagaggc caacccaaag atgccacgga ctgggtgctgc cagacccatg actgctgcta 240
 tgaccacctg aagacccagg ggtgcggcat ctacaaggac tattacagat acaacttttc 300
 ccaggggaac atccactgct ctgacaaggg aagctgggtg gagcagcagc tgtgtgctg 360
 tgacaaggag gtggccttct gcctgaagcg caacctggac acctaccaga agcgactgcg 420
 tttctactgg cggccccact gccgggggca gaccttggg tgctagaagc ccacaccctc 480
 taccctgttc ctcagcatgg agctctggca tccccacctc agtatctaac ctgaaccagc 540
 ctggcttttc aaacactcgg gggggaggta gtcccagcct ccccgggaac cc 592

<210> 5

<211> 2204

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> 28, 30

<223> a or g or c or t, unknown, or other

<220>

<221> misc_feature

<223> Incyte ID No: 1430683CB1

<400> 5

catggtttgg ggcgcagaga gggcaggan accaaggag agaagagggg aaattgcgcc 60
 cttttgggtg gaagctgtta tggtggacc ttaaagatc ttcgtagagt tgctgcccac 120
 cctggccctc tgtcttgaga gagggtctc tcacctcaca gacacaggat tattggctct 180
 tttctgcccc gccccctgcc cttttttttt tttttttgag atggagtctc tctctgtcgc 240
 ccaggctgga gtgcaatggc gtgatcttg ctcactgcaa cctccgcctc tggggttcaa 300
 gcgattctcc tgcctcagcc tcccagtag ctgggattac agactgaggg agctggccgt 360
 gcgactgggc ttcgggacct gtgcagagga gcaggcctc ctgagcagga ggaagcaggt 420
 ggtggccgcg gccttgagc aggcctgca gctggatgga gacctgcagg aggatgagat 480

```

cccagtggtta gctattatgg ccactgggtg tgggatccgg gcaatgactt ccctgtatgg 540
gcagctggctt ggcctgaagg agctgggctt cttggattgc gtctcctaca tcaccggggc 600
ctcgggctcc acctgggctt tggccaacct ttatgaggac ccagagtggc ctcagaagga 660
cctggcagggt cccactgagt tggtaagac ccaggtgacc aagaacaagc tgggtgtgct 720
ggcccccagc cagctgcagc ggtaccggca ggagctggcc gagcgtgccc gcttgggcta 780
cccaagctgc ttcaccaacc tgtgggacct catcaacgag gcgctgctgc atgatgagcc 840
ccatgatcac aagctctcag atcaacggga ggccttgagt catggccaga accctctgcc 900
catctactgt gccctcaaca ccaaagggca gagcctgacc acttttgaat ttggggagt 960
gtgcgagttc tctccctacg aggtcggtt ccccaagtac ggggccttca tccctctga 1020
gctctttggc tccgagttct ttatggggca gctgatgaag aggcttctg agtcccgcat 1080
ctgcttctta gaaggtatct ggagcaacct gtatgcagcc aacctccagg acagcttata 1140
ctgggacctca gagcccagcc agttctggga ccgctgggtc aggaaccagg ccaacctgga 1200
caaggagcag gtcccccttc tgaagataga agaaccaccc tcaacagccg gcagaatagc 1260
tgagtttttc accgatcttc tgacgtggcg tccactggcc caggccacac ataatttcct 1320
gcgtggcctc catttccaca aagactactt tcagcatcct cacttctcca catggaaagc 1380
taccactctg gatgggctcc ccaaccagct gacacctctg gagccccacc tgtgectgct 1440
ggatgttggc tacctcatca ataccagctg cctgccccct ctgcagccca ctcgggagct 1500
ggacctcatc ctgtcattgg actacaacct ccacggagcc ttccagcagt tgcagctcct 1560
gggccgggtc tgccaggagc aggggatccc gttcccaccc atctcgccca gccccgaaga 1620
gcagctccag cctcgggagt gccacacctt ctccgacccc acctgccccg gagccccctg 1680
gggtgctgac ttttctctg ggggtccggcg gacacccgag gaggcggcag ctggggaggt 1740
gaacctgtct tcatcggact ctccctacca ctacacgaag gtgacctaca gccaggagga 1800
cgtggacaag ctgctgcacc tgacacatta caatgtctgc aacaaccagg agcagctgct 1860
ggaggctctg cgccaggcag tgcagcggag gcggcagcgc aggccccact gatggccggg 1920
gccccctcca cccctaactc tcattcatct cctggctgct gagttgcagg tgggaactgt 1980
catcacgcag tgcttcagag cctcgggctc aggtggcact gtcccagggt ccaggctgag 2040
ggctgggagc tcccttgccg ctcagcagtt tgcagtgggg taaggaggcc aagcccattt 2100
gtgtaatcac ccaaaacccc ccggcctgtg cctgttttcc cttctgctgct accttgagta 2160
gttgaggcac ttgatacatc acagactcat aaaaaaaaaa aaaa 2204

```

<210> 6

<211> 1746

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1316804CB1

<400> 6

```

cagctctgag atttccagct cagcgatgcc cccaggctccc tgggagagct gcttctgggt 60
ggggggcctc attttgtggc tcagcggttg aagttcaggg gatgcacctc ctaccccaca 120
gccaaagtgc gctgacttcc agagcgccaa cctttttgaa ggcaccgatc tcaaagtcca 180
gtttctcttc tttgtccctt cgaatcctag ctgtgggcag ctagtagaag gaagcagtga 240
ctcccaaac tctgggttca atgccactct gggaacccaa ctaattatcc atggattcag 300
ggttttagga acaaagcctt cctggattga cacatttatt agaacccttc tgcgtgcaac 360
gaatgctaata gtgattgccg tggactggat ttatgggtct acaggagtct acttctcagc 420
tgtgaaaaat gtgattaagt tgagcctcga gatctccctt ttcctcaata aactcctggt 480
gctgggtgtg tcggaatcct caatccacat cattggtgtt agcctggggg cccacgttgg 540
gggcatggtg ggacagctct tcggaggcca gctgggacag atcacaggcc tggaccccg 600
tggacctgag tacaccaggg ccagtgtgga agagcgcttg gatgctggag atgccctctt 660
cgtggaagcc atccacacag acaccgacaa tttgggtatt cggattcccg ttggacatgt 720
ggactacttc gtcaacggag gccaaagacca acctggctgc cccaccttct ttacgcagg 780
ttatagttat ctgatctgtg atcacatgag ggtgtgtcac ctctacatca gcgccctgga 840

```

```

gaattcctgt ccactgatgg cctttccctg tgccagctac aaggccttcc ttgctggacg 900
ctgtctggat tgctttaacc cttttctgct ttcctgccca aggataggac tgggtggaaca 960
aggtggtgtc aagatagagc cgtcccccga ggaagtgaag gtctacctcc tgactacttc 1020
cagtgtcccg tactgcatgc atcacagcct cgtggagttt cacttgaagg aactgagaaa 1080
caaggacacc aacatcgagg ttaccttcct tagcagtaac atcacctctt catctaagat 1140
caccatacct aagcagcaac gctatgggaa aggaatcata gcccatgccca cccacacaatg 1200
ccagataaac caagtgaagt tcaagtttca gtcttccaac cgagtttggg aaaaagaccg 1260
gactaccatt attggaagt tctgcactgc ccttttgect gtcaatgaca gagaaaagat 1320
ggctctgctta cctgaaccag tgaacttaca agcaagtgtg actgtttcct gtgacctgaa 1380
gatagcctgt gtgtagttaa acctgggcag gacacatctc cctgcatttt tttttttttt 1440
tgagagagag gtgtgatgag ggatgtgtgt gtgcagctta ttgtagacca ttactactaa 1500
ggagaaaaag aaagctcttt cttattttcc tcataatcag ctaccctgga ggggagggag 1560
aactcatttt acagaacttg gtttcctttg ccgatcttat gtacataccc atttttagctt 1620
tcccatgcat acttaactgc acttgcttta tctccttggg cattcgact taggattcaa 1680
tagaaacatg tacagggtaa acaatttttt aaaaataaaa cttcatggag tatctgaaaa 1740
aaaaaa

```

<210> 7

<211> 146

<212> PRT

<213> Rattus norvegicus

<300>

<308> g204319

<400> 7

```

Met Lys Val Leu Leu Leu Leu Ala Val Val Ile Met Ala Phe Gly
  1                      5                      10                      15
Ser Ile Gln Val Gln Gly Ser Leu Leu Glu Phe Gly Gln Met Ile
                20                      25                      30
Leu Phe Lys Thr Gly Lys Arg Ala Asp Val Ser Tyr Gly Phe Tyr
                35                      40                      45
Gly Cys His Cys Gly Val Gly Gly Arg Gly Ser Pro Lys Asp Ala
                50                      55                      60
Thr Asp Trp Cys Cys Val Thr His Asp Cys Cys Tyr Asn Arg Leu
                65                      70                      75
Glu Lys Arg Gly Cys Gly Thr Lys Phe Leu Thr Tyr Lys Phe Ser
                80                      85                      90
Tyr Arg Gly Gly Gln Ile Ser Cys Ser Thr Asn Gln Asp Ser Cys
                95                      100                     105
Arg Lys Gln Leu Cys Gln Cys Asp Lys Ala Ala Ala Glu Cys Phe
                110                     115                     120
Ala Arg Asn Lys Lys Ser Tyr Ser Leu Lys Tyr Gln Phe Tyr Leu
                125                     130                     135
Asn Lys Phe Cys Lys Gly Lys Thr Pro Ser Cys
                140                     145

```

<210> 8

<211> 748

<212> PRT

<213> Gallus gallus

<300>

<308> g508625

<400> 8

```

Met Ser Phe Ile Asp Pro Tyr Gln His Ile Val Val Glu His Gln
  1           5           10           15
Tyr Ser His Val Phe Thr Val Thr Val Arg Lys Ala Thr Asn Val
          20           25           30
Thr Lys Gly Ala Ile Gly Asp Met Leu Asp Thr Pro Asp Pro Tyr
          35           40           45
Val Glu Leu Phe Ile Pro Ser Ala Pro Asp Cys Arg Lys Arg Thr
          50           55           60
Lys His Phe Asn Asn Asp Val Asn Pro Val Trp Asn Glu Thr Phe
          65           70           75
Glu Phe Ile Leu Asp Pro Asn Gln Asp Asn Val Leu Glu Val Thr
          80           85           90
Leu Met Asp Ala Asn Tyr Val Met Asp Glu Thr Leu Gly Met Ala
          95          100          105
Thr Phe Pro Ile Ser Ser Leu Lys Leu Gly Glu Lys Lys Glu Val
          110          115          120
Gln Leu Thr Phe Asn Asn Val Thr Glu Met Thr Leu Glu Leu Ser
          125          130          135
Leu Glu Val Cys Ser Ser Thr Asp Leu Arg Phe Ser Met Ala Leu
          140          145          150
Cys Asp Glu Glu Lys Lys Phe Arg Gln Gln Arg Lys Asp Asn Ile
          155          160          165
Met Gln Ser Met Lys Ser Phe Leu Gly Glu Glu Asn Ser Lys Asn
          170          175          180
Leu Thr Thr Ser Arg Asp Val Pro Val Ile Ala Val Leu Gly Ser
          185          190          195
Gly Gly Gly Phe Arg Ala Met Val Gly Phe Ala Gly Val Met Lys
          200          205          210
Ala Leu Tyr Glu Ser Gly Val Leu Asp Cys Ala Thr Tyr Ile Ala
          215          220          225
Gly Leu Ser Gly Ser Thr Trp Tyr Met Ser Thr Leu Tyr Ser His
          230          235          240
Pro Asp Phe Pro Glu Lys Gly Pro Lys Glu Ile Asn Gln Glu Leu
          245          250          255
Met Asn Ser Val Ser His Asn Pro Leu Leu Leu Leu Thr Pro Gln
          260          265          270
Lys Val Lys Arg Tyr Ile Glu Ala Leu Trp Asn Lys Lys Ser Ser
          275          280          285
Gly Gln Pro Val Thr Phe Thr Asp Ile Phe Gly Met Leu Ile Gly
          290          295          300
Glu Thr Leu Ile His Asn Arg Met Asp Thr Thr Leu Ser Asp Met
          305          310          315
Lys Glu Lys Val Ser Glu Ala Gln Cys Ala Leu Pro Leu Phe Thr
          320          325          330
Cys Leu His Val Lys Pro Asp Val Ser Glu Leu Met Phe Ala Asp
          335          340          345
Trp Val Glu Phe Ser Pro Tyr Glu Ile Gly Met Ala Lys Tyr Gly
          350          355          360
Thr Phe Met Ser Pro Asp Leu Phe Gly Ser Lys Phe Phe Met Gly
          365          370          375

```

Thr Val Val Lys	Lys Tyr Ser Glu Asn	Pro Leu His Phe Leu Met	
	380	385	390
Gly Val Trp Gly	Ser Ala Phe Ser Ile	Leu Phe Asn Arg Val Leu	
	395	400	405
Gly Val Ser Asn	Ser Gln Asn Lys Gly	Pro Thr Met Glu Glu Glu	
	410	415	420
Leu Glu Asn Ile	Arg Leu Lys His Leu	Val Ser Asn Asp Ser Ser	
	425	430	435
Asp Ser Glu Asp	Glu Ser Gln His Pro	Lys Gly Thr Glu Asn Ser	
	440	445	450
Glu Ala Asn Glu	Glu Tyr Gln Asn Ser	Ser Gln Glu Ser Trp Val	
	455	460	465
Gln Arg Met Leu	Met Ala Leu Val Gly	Asp Ser Ala Leu Phe Asn	
	470	475	480
Thr Arg Glu Gly	Arg Ala Gly Lys Val	His Asn Phe Met Leu Gly	
	485	490	495
Leu Asn Leu Asn	Ser Cys Tyr Pro Leu	Ser Pro Leu Ala Asp Leu	
	500	505	510
Leu Thr Gln Glu	Ser Val Glu Glu Asp	Glu Leu Asp Ala Ala Val	
	515	520	525
Ala Asp Pro Asp	Glu Phe Glu Arg Ile	Tyr Glu Pro Leu Asp Val	
	530	535	540
Lys Ser Lys Lys	Ile His Ile Val Asp	Ser Gly Leu Thr Phe Asn	
	545	550	555
Leu Pro Tyr Pro	Leu Ile Leu Arg Pro	Gln Arg Gly Val Asp Leu	
	560	565	570
Ile Ile Ser Phe	Asp Phe Ser Ala Arg	Pro Ser Asp Ser Ser Pro	
	575	580	585
Pro Phe Lys Glu	Ile Leu Leu Ala Glu	Lys Trp Ala Lys Met Asn	
	590	595	600
Lys Leu Pro Phe	Pro Lys Ile Asp Pro	Asn Val Phe Asp Arg Glu	
	605	610	615
Gly Leu Lys Glu	Cys Tyr Val Phe Lys	Pro Lys Asp Thr Ser Ser	
	620	625	630
Glu Lys Asp Cys	Pro Thr Ile Ile His	Phe Val Leu Ala Asn Ile	
	635	640	645
Asn Phe Arg Lys	Tyr Lys Ala Pro Gly	Leu Pro Arg Glu Ser Lys	
	650	655	660
Glu Glu Lys Asp	Phe Ala Asp Phe Asp	Ile Phe Asp Asp Pro Asn	
	665	670	675
Thr Pro Phe Ser	Thr Phe Asn Phe Gln	Tyr Pro Asn Glu Ala Phe	
	680	685	690
Lys Arg Leu His	Asp Leu Met Glu Phe	Asn Thr Leu Asn Asn Leu	
	695	700	705
Asp Val Ile Lys	Gln Ala Met Met Glu	Ser Ile Glu Tyr Arg Lys	
	710	715	720
Glu Asn Pro Ser	Arg Cys Ser Val Ser	Leu Ser Ser Val Glu Ala	
	725	730	735
Arg Arg Phe Phe	Asn Lys Asn Asn Leu	Asn Asn His Thr	
	740	745	

<210> 9
 <211> 456
 <212> PRT

<400> 9														
Met	Cys	Pro	Gly	Leu	Trp	Gly	Thr	Cys	Phe	Trp	Leu	Trp	Gly	Ser
1				5					10					15
Leu	Leu	Trp	Leu	Ser	Ile	Gly	Arg	Ser	Gly	Asn	Val	Pro	Pro	Thr
				20					25					30
Thr	Gln	Pro	Lys	Cys	Thr	Asp	Phe	Gln	Ser	Ala	Asn	Leu	Leu	Arg
				35					40					45
Gly	Thr	Asn	Leu	Lys	Val	Gln	Phe	Leu	Leu	Phe	Thr	Pro	Ser	Asp
				50					55					60
Pro	Gly	Cys	Gly	Gln	Leu	Val	Glu	Glu	Asp	Ser	Asp	Ile	Arg	Asn
				65					70					75
Ser	Glu	Phe	Asn	Ala	Ser	Leu	Gly	Thr	Lys	Leu	Ile	Ile	His	Gly
				80					85					90
Phe	Arg	Ala	Leu	Gly	Thr	Lys	Pro	Ser	Trp	Ile	Asn	Lys	Phe	Ile
				95					100					105
Arg	Ala	Leu	Leu	Arg	Ala	Ala	Asp	Ala	Asn	Val	Ile	Ala	Val	Asp
				110					115					120
Trp	Val	Tyr	Gly	Ser	Thr	Gly	Met	Tyr	Phe	Ser	Ala	Val	Glu	Asn
				125					130					135
Val	Val	Lys	Leu	Ser	Leu	Glu	Ile	Ser	Arg	Phe	Leu	Ser	Lys	Leu
				140					145					150
Leu	Glu	Leu	Gly	Val	Ser	Glu	Ser	Ser	Ile	His	Ile	Ile	Gly	Val
				155					160					165
Ser	Leu	Gly	Ala	His	Val	Gly	Gly	Met	Val	Gly	His	Phe	Tyr	Lys
				170					175					180
Gly	Gln	Leu	Gly	Arg	Ile	Thr	Gly	Leu	Asp	Pro	Ala	Gly	Pro	Glu
				185					190					195
Tyr	Thr	Arg	Ala	Ser	Leu	Glu	Glu	Arg	Leu	Asp	Ser	Gly	Asp	Ala
				200					205					210
Leu	Phe	Val	Glu	Ala	Ile	His	Thr	Asp	Thr	Asp	Asn	Leu	Gly	Ile
				215					220					225
Arg	Ile	Pro	Val	Gly	His	Val	Asp	Tyr	Phe	Val	Asn	Gly	Gly	Gln
				230					235					240
Asp	Gln	Pro	Gly	Cys	Pro	Ala	Phe	Ile	His	Ala	Gly	Tyr	Ser	Tyr
				245					250					255
Leu	Ile	Cys	Asp	His	Met	Arg	Ala	Val	His	Leu	Tyr	Ile	Ser	Ala
				260					265					270
Leu	Glu	Asn	Thr	Cys	Pro	Leu	Met	Ala	Phe	Pro	Cys	Ala	Ser	Tyr
				275					280					285
Lys	Ala	Phe	Leu	Ala	Gly	Asp	Cys	Leu	Asp	Cys	Phe	Asn	Pro	Phe
				290					295					300
Leu	Leu	Ser	Cys	Pro	Arg	Ile	Gly	Leu	Val	Glu	Arg	Gly	Gly	Val
				305					310					315
Lys	Ile	Glu	Pro	Leu	Pro	Lys	Glu	Val	Arg	Val	Tyr	Leu	Gln	Thr
				320					325					330
Thr	Ser	Ser	Ala	Pro	Tyr	Cys	Val	His						

Phe	Leu	Gly	Asn	Asn	Val	Thr	Ser	Ser	Val	Lys	Ile	Thr	Ile	Pro	
				365						370				375	
Lys	Asp	His	Leu	Glu	Gly	Arg	Gly	Ile	Ile	Ala	His	Gln	Asn	Pro	
				380						385				390	
His	Cys	Gln	Ile	Asn	Gln	Val	Lys	Leu	Lys	Phe	His	Ile	Ser	Ser	
				395						400				405	
Arg	Val	Trp	Arg	Lys	Asp	Arg	Thr	Pro	Ile	Val	Gly	Thr	Phe	Cys	
				410						415				420	
Thr	Ala	Pro	Leu	Pro	Val	Asn	Asp	Ser	Lys	Lys	Thr	Val	Cys	Ile	
				425						430				435	
Pro	Glu	Pro	Val	Arg	Leu	Gln	Val	Ser	Met	Ala	Val	Leu	Arg	Asp	
				440						445				450	
Leu	Lys	Met	Ala	Cys	Val										
				455											

<210> 10

<211> 5

<212> PRT

<213> Homo sapiens

<220>

<221> unsure

<222> 2, 4

<223> unknown, or other

<220>

<221> misc_feature

<223> motif

<400> 10

Gly Xaa Ser Xaa Gly

1

5